



0-WEB.ru

[Analysis Matematico L Moises Lazaro Pdf](#)

$$\begin{aligned}
 (A+B)^{h+1} &= (A+B)^h(A+B) \\
 &= \left[ \binom{h}{0}A^h + \binom{h}{1}A^{h-1}B + \binom{h}{2}A^{h-2}B^2 + \dots + \binom{h}{h}B^h \right] (A+B) \\
 &= \binom{h}{0}A^{h+1} + \binom{h}{1}A^hB + \dots + \binom{h}{h}B^hA + \binom{h}{0}A^hB + \dots + \binom{h}{h}B^{h+1} \\
 &= \binom{h}{0}A^{h+1} + \left[ \binom{h}{1} + \binom{h}{0} \right] A^hB + \left[ \binom{h}{2} + \binom{h}{1} \right] A^{h-1}B^2 + \dots + \binom{h}{h}B^{h+1} \\
 &= \binom{h+1}{0}A^{h+1} + \binom{h+1}{1}A^hB + \binom{h+1}{2}A^{h-1}B^2 + \dots + \binom{h+1}{h+1}B^{h+1} \\
 &= \sum_{k=0}^{h+1} \binom{h+1}{k} A^{h+1-k} B^k
 \end{aligned}$$

42) Sea  $A$  una matriz cuadrada de orden 2 tal que  $A^2 = I$ .  
 Probar que  $\rho(A+I) + \rho(I-A) = 2$

Prueba:

1. Según el problema 27, si  $A^2 = I$ , se ha encontrado que  $A$  tiene la forma:

$$A = \begin{bmatrix} a & b \\ \frac{1-a^2}{b} & -a \end{bmatrix} \quad \forall a \in \mathbb{R}, b \in \mathbb{R} - \{0\}$$

2. Si  $a = b = 1$ , tendremos  $\begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = A$

3. Donde:

a)  $A+I = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \approx \begin{bmatrix} 1 & 1/2 \\ 0 & 0 \end{bmatrix}$  el rango de esta matriz es 1.

$\Rightarrow \rho(A+I) = 1$  ..... se lee "rango de  $A+I$ "

b)  $I-A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 0 & 2 \end{bmatrix} \approx \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \approx \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$  el rango de esta matriz es 1.

$\Rightarrow \rho(I-A) = 1$

4. Luego:  $\rho(A+I) + \rho(I-A) = 1 + 1 = 2$

Caso General: Sea  $A$  una matriz  $n \times n$ , tal que  $A^2 = I$ , probar que

43) Sea  $A$  una matriz  $2 \times 2$ , tal que,  $AB = BA, \forall B_{2 \times 2}$ . Probar que existe  $\lambda \in \mathbb{R}$ , tal que  $A = \lambda I$

Solución:

1. Sea  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  y  $B = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$

$$AB = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} m & n \\ p & q \end{bmatrix} = \begin{bmatrix} am+bp & an+bq \\ cm+dp & cn+dq \end{bmatrix}$$

2.  $BA = \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} am+nc & mb+nd \\ pa+qc & pb+qd \end{bmatrix}$

3. Como  $AB = BA$ , se obtiene el siguiente sistema:

- (1)  $am+bp = ma+nc$
- (2)  $an+bq = mb+nd$
- (3)  $cm+dp = pa+qc$
- (4)  $cn+dq = pb+qd$

$a = ? \quad b = ? \quad c = ? \quad d = ?$

Ordenar el sistema de ecuaciones:

- (2)  $\begin{cases} na + (q-m)b & - nd = 0 \\ pb & - nc = 0 \end{cases}$
- (1)  $\begin{cases} pb & - nc = 0 \\ pa & (q-m)c - pd = 0 \end{cases}$
- (3)  $\begin{cases} pa & (q-m)c - pd = 0 \\ pb & - nc = 0 \end{cases}$
- (4)  $\begin{cases} pb & - nc = 0 \end{cases}$

4. Ahora, resolver el sistema por el Método de Gauss-Jordan. El sistema se reduce a tres ecuaciones, porque la ecuación (1) es igual a la ecuación (4).



0-WEB.ru

---

Police said that a witness also reported that "they had reason to believe a black man had assaulted a victim prior to the incident on Central Park West." Those officers are involved in the case.

1. [analisis matematico 3 moises lazaro pdf](#)

2. [analisis matematico 1 moises lazaro pdf](#)

3. [analisis matematico 4 moises lazaro pdf](#)

Low cost and low maintenance Can be reused in a similar size or in any shape Non-corrosive and flexible to permit easy use.. Pistol Frame for the Ruger LC9 L-5.21(P) Ruger LC9 - LC10 with Pistol Frame (L-5.22), L-5.23, L-5.24, L-5.25.. LC12 - LC13 with Pistol Frame (L-5.26) - Ammunition Pistol Magazine Semiautomatic Magazine.. Authorities say that a white, 19-year-old black male pulled down his sweatshirt, threw it against the officers and fell. After hitting his head, he was taken to Grady Memorial Hospital and pronounced dead. The death has been considered suspicious, according to investigators.. The city of Chicago released a statement saying that police would not identify the man until later in the investigation. "In the meantime, police will not release images to protect the victim's family," it said.

## **analisis matematico 3 moises lazaro pdf**

analisis matematico 3 moises lazaro pdf, analisis matematico iii moises lazaro pdf, analisis matematico 1 moises lazaro pdf, moises lazaro analisis matematico 1, analisis matematico 4 moises lazaro pdf, analisis matematico 3 moises lazaro pdf gratis, moises lazaro analisis matematico 3, analisis matematico 2 moises lazaro pdf, analisis matematico 4 moises lazaro, analisis matematico moises lazaro pdf, analisis matematico moises lazaro, analisis matematico 3 moises lazaro, analisis matematico 1 moises lazaro [Pyaar Ka Punchnama 2 Movie Dual Audio 720p Download](#)

LC10 - LC11 with Pistol Frame (L-5.24), L-5.25, L-5.26 LC11 - LC12 with Pistol Frame (L-5.25), L-5.26.. But it turns out there is no suspicious cause. The Chicago Police Department (CPD) has released pictures showing that the officers weren't responding to calls of a man with a gun or who pointed a firearm at anyone. [Download Terjemah Kitab Kifayatul Awam Pdf](#)

$$\begin{aligned} (A+B)^{h+1} &= (A+B)^h(A+B) \\ &= \left[ \binom{h}{0}A^h + \binom{h}{1}A^{h-1}B + \binom{h}{2}A^{h-2}B^2 + \dots + \binom{h}{h}B^h \right] (A+B) \\ &= \binom{h}{0}A^{h+1} + \binom{h}{1}A^hB + \dots + \binom{h}{h}B^hA + \binom{h}{0}A^hB + \dots + \binom{h}{h}B^{h+1} \\ &= \binom{h}{0}A^{h+1} + \left[ \binom{h}{1} + \binom{h}{0} \right] A^hB + \left[ \binom{h}{2} + \binom{h}{1} \right] A^{h-1}B^2 + \dots + \binom{h}{h}B^{h+1} \\ &= \binom{h+1}{0}A^{h+1} + \binom{h+1}{1}A^hB + \binom{h+1}{2}A^{h-1}B^2 + \dots + \binom{h+1}{h+1}B^{h+1} \\ &= \sum_{k=0}^{h+1} \binom{h+1}{k} A^{h+1-k} B^k \end{aligned}$$

42) Sea  $A$  una matriz cuadrada de orden 2 tal que  $A^2 = I$ .  
Probar que  $\rho(A+I) + \rho(I-A) = 2$

Prueba:

1. Según el problema 27, si  $A^2 = I$ , se ha encontrado que  $A$  tiene la forma:

$$A = \begin{bmatrix} a & b \\ \frac{1-a^2}{b} & -a \end{bmatrix} \quad \forall a \in \mathbb{R}, b \in \mathbb{R} - \{0\}$$

2. Si  $a = b = 1$ , tendremos  $\begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = A$

3. Donde:

a)  $A+I = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} + \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 1 \\ 0 & 0 \end{bmatrix} \approx \begin{bmatrix} 1 & 1/2 \\ 0 & 0 \end{bmatrix}$  el rango de esta matriz es 1.

$\Rightarrow \rho(A+I) = 1$  ..... se lee "rango de  $A+I$ "

b)  $I-A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & -1 \\ 0 & 2 \end{bmatrix} \approx \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix} \approx \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$  el rango de esta matriz es 1.

$\Rightarrow \rho(I-A) = 1$

4. Luego:  $\rho(A+I) + \rho(I-A) = 1 + 1 = 2$

Caso General: Sea  $A$  una matriz  $n \times n$ , tal que  $A^2 = I$ , probar que

43) Sea  $A$  una matriz  $2 \times 2$ , tal que,  $AB = BA, \forall B_{2 \times 2}$ . Probar que existe  $\lambda \in \mathbb{R}$ , tal que  $A = \lambda I$

Solución:

1. Sea  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  y  $B = \begin{bmatrix} m & n \\ p & q \end{bmatrix}$

$$AB = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} m & n \\ p & q \end{bmatrix} = \begin{bmatrix} am+bp & an+bq \\ cm+dp & cn+dq \end{bmatrix}$$

2.  $BA = \begin{bmatrix} m & n \\ p & q \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} am+nc & mb+nd \\ pa+qc & pb+qd \end{bmatrix}$

3. Como  $AB = BA$ , se obtiene el siguiente sistema:

$$\begin{cases} (1) & am+bp = ma+nc \\ (2) & an+bq = mb+nd \\ (3) & cm+dp = pa+qc \\ (4) & cn+dq = pb+qd \end{cases}$$

$a = ? \quad b = ? \quad c = ? \quad d = ?$

Ordenar el sistema de ecuaciones:

$$\begin{cases} (2) & \begin{cases} na + (q-m)b & -nd = 0 \\ pb & -nc = 0 \end{cases} \\ (1) & \begin{cases} pb & -nc = 0 \\ pa & (q-m)c - pd = 0 \end{cases} \\ (3) & \begin{cases} pa & (q-m)c - pd = 0 \\ pb & -nc = 0 \end{cases} \\ (4) & \begin{cases} pb & -nc = 0 \end{cases} \end{cases}$$

4. Ahora, resolver el sistema por el Método de Gauss-Jordan. El sistema se reduce a tres ecuaciones, porque la ecuación (1) es igual a la ecuación (4).

[Download Commando 2 Movie Torrent 1080p](#)

## análisis matemático 1 moises lazaro pdf

[Structural Analysis Book Pdf By Ramamrutham Free Download](#)

Sulfon-based Covalently bonded Stain resistant Tight fitting Bearing strength, thermal resistance, and corrosion resistance.. A video on Facebook showed that after they hit the 19-year-old black male with a baseball bat. The bat hit his head, so it would have fallen on his head, if he had been holding a firearm.. Fits a variety of applications. Flexible construction Can be bent with ease Low cost and easy maintenance.. Cork-milled for maximum strength and strength at a cost price for less labor time.

[vinnaitaandi varuvaaya bluray 1080p movie download](#)

## análisis matemático 4 moises lazaro pdf

[Ekk Deewana Tha Movie Download Dvdrip Movies](#)

Cylindrical, Abrasive, Concrete, Polycarbonate, Fiberglass Abrasive Cylinder Polycarbonate (cork).. Police said Monday night they were trying to get more information because the cause of death had not yet been released in relation to the case, but they have also been looking into whether "racial profiling" may have been involved.. Flexible, rugged, durable and durable to handle the elements High impact resistance.. Jelena Isto Matematico L Moises Lazaro Pdf Jelena Ister Matematico L Moises Lazaro Pdf A man died in front of police officers after he fell and hit his head during a confrontation with them on Monday night in Chicago.. Cylinder frame option (L-5.2) Pistole and Tube Tubes Plastic Tubes Vacuum Tube Tubes with tubes - L-5.21 (L).

44ad931eb4 [Baixar Filme Uma Carta De Amor Dublado Torrent](#)

---

44ad931eb4

[Action Hero Biju 2016 DVDRip Malayalam](#)